



Age and Sex Referents Quiz by Laura King, MA, ELS

Directions: Edit the following sentences to eliminate incorrect age and sex referents. Refer to [section 11.5](#) (p 410 in print) of the [AMA Manual of Style](#).

1. A subpopulation at higher risk for sexually transmitted disease may be African American adolescent females seeking sexual health services.
2. The study participants ranged in age from 10 to 25 years and included both boys and girls.
3. Incidence was estimated as the proportion of newborn babies meeting the specified cutoff for mild hypothermia or for moderate or severe hypothermia.
4. The infants were examined immediately after birth for the presence of hypoxic-ischemic encephalopathy.

LEARNING RESOURCES

5. The investigators studied the incidence of acute respiratory illness in children 1 to 12 months old.

6. Patients included in the study had a mean age of 6½ years (range, 2-15 years).

7. The study examined potential major bleeding events in the elderly (those older than 75 years).

8. The study examined the risk of human immunodeficiency virus among males and females treated at the local community health clinic.

9. From the population sample, 25 men (mean age, 35 years; range, 26-49 years) and 25 girls (mean age, 32 years; range, 24-47 years) were selected for participation in the study.

10. The primary outcome of the study was hearing assessment of adolescents 18 to 24 years of age with various types of craniosynostosis.

